

Magnetically Driven Outflows in a Starburst Environment

Elisabete M. de Gouveia Dal Pino^{1,2} & Gustavo A. Medina Tanco^{1,3}

Received _____; accepted _____

¹Instituto Astronômico e Geofísico, University of São Paulo, Av. Miguel Stéfano, 4200, São Paulo 04301-904, SP, Brasil; E-mail: dalpino@iagusp.usp.br, gustavo@iagusp.usp.br

² Astronomy Department, Theoretical Astrophysics Center, University of Berkeley, 601 Campbell Hall, Berkeley, CA 94720-3411

³Department of Physics and Astronomy, University of Leeds, EC Stoner Building, Leeds LS2 9JT, United Kingdom

ABSTRACT

We here investigate the possibility that the observed collimated outflows in luminous infrared galaxies (LIGs) and some Seyfert galaxies can be produced in a starburst (SB) environment. In the former source class, in particular, there seems to be some observational evidence for the presence of nuclear SBs in some objects (e.g. Scoville, Lonsdale, & Lonsdale 1998).

A nuclear disk can be quickly produced by gas infall during star formation in a *rotating*, stellar cluster. We find that massive nuclear SBs with core disk masses $M_d \sim 10^8 - 10^9 M_\odot$, and supernova rates $\nu_{SN} \simeq 5 \times 10^{-3} - 2 \text{ yr}^{-1}$ (which are consistent with the ν_{SN} values inferred from the observed non-thermal radio power in source candidates) may inject kinetic energies which are high enough to blow out directed flows from the accreting disk surface, within the SB lifetimes. In our models, the acceleration and collimation of the nuclear outflow are provided by magnetic fields anchored into the rotating SB-disk. The emerging outflow carries a kinetic power that is only a small fraction (a few percent) of the supernovae energy rate produced in the SB. Based on conditions determined from observed outflows and disks, we find that moderate disk magnetic fields ($\gtrsim 8 \times 10^{-4} \text{ G}$) are able to accelerate the outflows up to the observed terminal velocities ($\lesssim \text{few } 100 \text{ km s}^{-1}$ in the case of the Seyfert galaxies, and $\sim 400 - 950 \text{ km s}^{-1}$ in the case of the LIGs). The outflow is produced within a wind zone in the disk of radius $\lesssim 100 \text{ pc}$ in the the LIGs, and $\lesssim 10 \text{ pc}$ in the Seyferts, with wind mass loss to disk accretion rate ratios $\dot{M}_w/\dot{M}_d \gtrsim 0.1$ (where $\dot{M}_d \sim 100 M_\odot \text{ yr}^{-1}$). The observation of rotating nuclear disks of gas within $\sim \text{few } 100 \text{ pc}$ scales, like that in the prototype LIG, Arp 220 (for which $\dot{M}_d \sim 100 M_\odot \text{ yr}^{-1}$), and magnetized outflows in Sey galaxies and LIGs (with terminal $B \sim 10^{-5} \text{ G}$ at the kpc-scales), provide some observational

support for the magneto-centrifugal disk picture drawn here.

Subject headings: **galaxies: active - ISM: jets and outflows - galaxies:
starburst, Seyfert galaxies, LIGs**

1. Introduction

Active Galactic Nuclei (AGN) emit large amounts of energy (10^{43-48} ergs s^{-1} , from Seyfert galaxies to QSOs) over a large range of the spectrum coming from a small volume. The nature of the central engine that produces these extraordinary amounts of energy remains virtually unknown. Current AGN theories include: i) black holes (BHs) accreting matter from the surroundings (e.g. Rees 1984); ii) Starburst (SB) models in which activity is due to star formation in a central cluster (e.g. Terlevich et al. 1992 - who have revived old theories, e.g., of Shklovskii 1960, Field 1964); and iii) BHs surrounded by dense starbursts (e.g. Perry & Dyson 1985, Scoville & Norman 1988, Williams & Perry 1994).

While BH models (with masses $\sim 10^6 - 10^{10} M_{\odot}$) accreting matter from the surroundings have the power to produce the observed luminosities and naturally provide a preferred axis for collimating jets, they seem to explain unsatisfactorily, e.g., the broad emission-line regions (BELRs) of AGN (as they require a fine-tuning to fit the observations). In Starburst models, on the other hand, strongly radiative compact supernova remnants (cSNRs) are the source of the broad emission lines. This model has been proved plausible in explaining the observed emission line spectrum, the UV-optical continuum and related variability (e.g., Terlevich et al. 1992), but is difficult to be applied to the whole population of AGN. In particular, they do not provide a natural mechanism for jet production and collimation. We here address this question.

It seems plausible that active star formation and a compact AGN activity may coexist in the nucleus of luminous infrared galaxies (the so called LIGs which are characterized by extreme mid- to far-IR luminosities $L > 10^{11} L_{\odot}$ and high brightness temperature radio emission). The question to assess the relative contribution of each form of activity to the LIGs energetics, and to determine a potential evolutionary progression from one to the other has been recently addressed by Smith, Lonsdale & Lonsdale (1998, hereafter SLL98).

Searching for a SB-compact object AGN connection in these luminous infrared galaxies they have demonstrated that nuclear starbursts can in principle explain the observed VLBI- and VLA- scale structures and the high infrared luminosities of most of these objects, although in some cases severe constraints must be imposed on the initial mass function and, in general, complexes of extremely luminous radio supernovae (RSNs) (clustered in pc-scale clumps) are required to explain the high- T_b observed. In a sample of 10 LIGs, they found potentially 6 SB systems, 3 possible hybrid/intermediate systems and 1 quasar-like object (or BH candidate). This would favor an evolutionary interpretation in which a SB eventually begins to fuel a coalescing central compact object. Galactic mergers are believed to be responsible for the origin of LIGs, and numerical simulations (Mihos & Hernquist 1996) suggest that the SB occurs relatively late in the merging process ($\sim 10^8$ yr). SLL98 results, in turn, suggest that a SB phase might carry on for another $\sim 10^8$ yr as the nuclei coalesce and a compact AGN develops. Once the compact AGN fully turns on, it must quickly consume its shroud of dust thus extinguishing the star formation process.

A prototype of the LIG class, Arp 220, seems to be dominated by a massive nuclear starburst imbedded in a rotating disk-like structure of dense molecular gas with a radius ~ 235 pc, and with densities ($n \sim 10^4$ cm $^{-3}$) and temperatures similar to those of the GMC core regions in our Galaxy (Scoville, Yun, & Bryant 1997, hereafter SYB97). Several other systems (e.g., NGC 6240 and Mrk231) bear striking similarities to Arp 220, which are suggestive of a centrifugally supported disk of dense molecular gas. These disks may play a critical role in the evolution of ultra-luminous galactic nuclei and in their energy release. The high density of the gas will undoubtedly promote extremely high rates of massive star formation since the conditions are very similar to those of GMC cores. In these cases, the presence of strong, highly ordered magnetic fields could provide both the conditions appropriate for RSN formation and production of collimated outflows.

Nuclear outflows have been observed in several Seyfert galaxies and LIGs (e.g., Arp 220, NGC 6240, NGC 2782, NGC 4945, and NGC 4235) (see, e.g., Heckman, Lehrnet, & Armus 1993, Colbert et al. 1996, Joglee et al. 1998, and references therein). In the case of the LIGs, the detection of emission line profiles with typical line widths of 200-600 km s⁻¹, and in some cases with FWHM as large as 1000 km s⁻¹ (e.g. NGC 6240), are signatures of the presence of high-speed, supersonic outflows where the lines are produced by shock-heated ambient gas which is swept by the outflow (Heckman, Armus, & Miley 1990, hereafter HAM90). In the case of the Seyfert galaxies, radio observations of a distance-limited sample of edge-on galaxies by Colbert et al. (1996), have shown that 6 of the 10 observed galaxies, have extended radio structures with radial extent between 5 and 30 kpc. They are oriented at skewed angles ($\Delta = 35^\circ - 90^\circ$) with respect to the galaxy minor axis and do not look like the wide-angled winds observed in typical starburst galaxies. Their morphologies resemble instead, those of more collimated outflows originating in the nuclear region (within 1 kpc), like the jets observed in extended radio galaxies, although the Seyfert outflows are weaker (with radio power at 4.9 GHz, $P_{4.9} = 10^{20-22}$ W Hz⁻¹), smaller, and less collimated than those of the radio galaxies (which have typical radio power at 1.4 GHz, $P_{1.4} > 10^{24}$ W Hz⁻¹). Terminal outflow velocities of ~ 25 to 150 km s⁻¹ and magnetic fields $\sim 10^{-7}$ to 10^{-6} G at 5 to 30 kpc scales (implying average fields $\sim 10^{-5}$ G at ~ 1 kpc) have been inferred from the observations.

Based on the evidences above, it seems plausible that the LIGs and perhaps some of the low-level-activity AGN class of Seyfert galaxies may be powered by nuclear SBs at least during part of their active lives. In the present work, we assume that a SB, which has been induced in an interaction or merging process, is in progress in the nuclear region of these objects and examine the conditions under which collimated outflows can be produced in such a scenario. Previous purely hydrodynamic numerical studies of starbursts generated within few kpc radius from the nucleus have shown that SB mass loss dynamics may

provide favorable conditions for a free vertically directed escape of hot gas from the galactic environment at the vigorous SN dominated phase of the SB (e.g., Tomisaka & Ikeuchi 1988, De Young & Gallagher 1990, Suchkov et al. 1994, Tenorio-Tagle & Muñoz-Tuñón 1997). Those models have successfully explained the formation of the poor-collimated, wide-angled super-winds (with radius ~ 1 kpc) often observed in SB galaxies (e.g., HAM90). In this work, however, we are mainly interested in systems with potential SB-AGN activity connection and thus examine the production of outflows within radial distances of few tens of parsecs from the central core. We consider a mechanism that can provide efficient collimation and acceleration of the emerging outflow on these scales. We assume that part of the gas released by star formation in the rotating SB system is quickly collected into a disk with similar conditions, for example, to those observed in the Arp 220 nuclear disk. Observations support the evidence for magnetized outflows. Since magnetic winds can be produced by rotating objects with a magnetic field anchored into them (e.g., Weber and Davis 1967), we assume that magnetic fields are present due to magnetic-flux captured from the environment during the formation of the disk, and compression by the accretion process (see, e.g., Spruit 1996).

The perception that rapid rotation coupled with strong magnetic fields could drive outflows was first realized in the context of the solar wind theory (e.g., Weber and Davis 1967, Mestel 1968). Later, Blandford & Payne (1982) introduced a scale free hydromagnetic wind model for jet acceleration from a magnetized disk which gives centrifugally driven outflow solutions. These solutions also have the possibility of being self-collimated (Heyvaerts & Norman 1989). (See, however, Okamoto 1998 for a discussion on the limitations of these models.) Similar outflow models were subsequently applied to proto-stellar objects (e.g., Pudritz & Norman 1983, Pelletier & Pudritz 1992, Wardle & Königl 1993, Shu et al. 1994, see also Spruit 1996, for a review), galactic X-ray binaries (e.g., Blandford 1993), and proto-jovian planets (Quillen & Trillen 1998). In the present

work, we use the Pelletier and Pudritz solutions (hereafter PP92) to investigate the formation of magneto-centrifugally accelerated outflows in a SB environment.

In §2 we try to build a consistent SB scenario and examine whether massive nuclear star formation may be present at high enough rates to drive galactic flows out from the surface of the nuclear gas disk produced in the course of the SB evolution. In §3, we discuss the physical conditions for the nuclear disk to produce collimated outflows based on the observed characteristics of the Arp 220 nuclear disk, and estimate the radius, accretion and wind outflow rates, terminal velocities, kinetic energy rates, and magnetic field ranges for which magneto-centrifugal outflows could operate. In §4, we draw our conclusions and briefly discuss our results.

2. A Starburst Scenario

In an attempt to build consistent starburst scenarios and evaluate their energetics, mass loss, and rate of supernova production, in previous work (de Gouveia Dal Pino & Medina Tanco 1997, hereafter GT97) we have calculated the evolution of several starburst stellar clusters based on the evolutionary population synthesis models of Schaller et al. (1992). As initial conditions we have assumed a Salpeter initial stellar mass distribution function [$\Phi(m) \propto m^{-\gamma}$], with a spectral index $\gamma = 2.35$, and stellar masses in the range [$M_l = 1 - 8 M_\odot$, $M_u = 120 M_\odot$]; initial number of stars $N_c = 10^6 - 4 \times 10^9$ (corresponding to cluster radii $R_c \simeq 0.2 - 705$ pc and masses $M_c \simeq 10^7 - 10^{10} M_\odot$); and an energy deposited by each supernova (SN) $E = 10^{51}$ erg s⁻¹. Assuming also that the system was rotating with an ellipticity $\epsilon = J/r_c v_s \simeq 2.5 \times 10^{-3}$, where J is the angular momentum per unit mass (Salpeter and Saslaw 1996, Begelman & Rees 1978), we have found that part of the gas released by star formation may cool quickly by free-free emission in time scales much smaller than the evolutionary time scales of the system ($\tau \lesssim 10^8$ yr) and thus fall towards

the centre and settle into a disk with an initial characteristic radius $r_d \simeq \epsilon^{1/2} r_c$. In fact, rapid fueling is expected by the large star formation rates and gas conversion into stars detected in nuclear SB regions (Kennicutt et al. 1987, Lo et al. 1987). These observations indicate that fueling should occur in time scales smaller than the conversion time into stars ($\sim 10^7 - 10^8$ yr) if the gas is to accumulate faster than it is consumed (e.g., Tenorio-Tagle & Muñoz-Tuñón 1997). As an example, Figure 1 (see also GT97) shows the results for a cluster with $3 \times 10^8 M_\odot$ and a core disk which eventually becomes also gravitationally unstable and undergoes star formation (see below). For a cluster-disk system with masses and radii about ten times larger (smaller) than those of Fig. 1, all the luminosity values and supernova rates increase (decrease) by an order of magnitude.

As investigated in previous SB-driven outflow models, a strong nuclear SB can, in principle, drive a wide-angled wind out of the system (see e.g., HAM90). We will first discuss this process in terms of the energy available to produce winds in a SB system and then, on subsequent sections, examine the mechanism for providing outflow acceleration and collimation. If an outflow is driven by a SB, then the observed non-thermal radio power (P_{nt}) should be correlated with the supernovae production rate (ν_{SN}) in the SB through (Condon & Yin 1990):

$$P_{nt} \simeq 1.3 \times 10^{23} \text{ WHz}^{-1} \nu_{SN} \left(\frac{\nu}{1 \text{ GHz}} \right)^{-\alpha_r} \quad (1)$$

where ν is the observed radio frequency, and $\alpha_r \simeq 0.7 - 0.8$ is the spectral index. Using the observed radio power for the Seyfert outflows we find (see Condon et al. 1996) $\nu_{SN} \simeq 5.1 \times (10^{-3} - 10^{-1}) \text{ yr}^{-1}$, which is consistent with the predicted values of ν_{SN} by the SB models (see, e.g., Fig. 1).

Alternatively, simple analytical prescription can be used to estimate the expected supernova production rate ν_{SN} as a function of the star formation rate (SFR). For a power law initial stellar mass function, with upper and lower stellar mass cutoffs $m_u \simeq 120 M_\odot$,

and $m_l \simeq 1M_\odot$, respectively, we have (e.g., Scoville & Soifer 1991)

$$\nu_{SN} = \int \Phi(m) dm \simeq \frac{SFR}{3} \frac{(m_{SN}^{-1.35} - m_u^{-1.35})}{(m_l^{-0.35} - m_u^{-0.35})} \quad (2)$$

where $m_{SN} = 8M_\odot$ is the minimum mass for supernova detonation, or

$$\nu_{SN} \simeq 0.5 \text{ yr}^{-1} \left(\frac{SFR}{20 M_\odot \text{ yr}^{-1}} \right) \quad (3)$$

which shows that $SFR \simeq (5 - 20) M_\odot \text{ yr}^{-1}$ result SN rates consistent with those required to explain the observed radio power in Sey outflows.

In the case of the LIGs, the observed infrared luminosities imply star formation rates $SFR \simeq 16 - 125 M_\odot \text{ yr}^{-1}$. SBs models with these SFR result in supernova rates $\nu_{SN} \simeq 0.1 - 2 \text{ yr}^{-1}$ (SLL98). These are also consistent with the ν_{SN} values required by the observed non-thermal radio luminosity in these objects. In the particular case of the Arp 220 source, the supernova rate required by the observed non-thermal radio power is $\nu_{SN} \simeq 2 \text{ yr}^{-1}$ (which is in agreement with the predicted rate by a SB model with cluster and disk masses an order of magnitude larger than those shown in Fig. 1, $M_d \simeq 10^9 M_\odot$).

SBs with $\nu_{SN} \simeq 5 \times 10^{-3} - 2 \text{ yr}^{-1}$ may inject *kinetic* energy into the ambient medium at a rate (see, e.g., Eq. 1; HAM90)

$$L_K \simeq 3.5 \times 10^{43} \nu_{SN} \simeq (2 \times 10^{41} - 7 \times 10^{43}) \text{ ergs}^{-1} \quad (4)$$

which is comparable to both the rate of gravitational energy released by the infalling gas and the SN luminosity in the core disk of the SB models (see, e.g., Fig. 1, for which $\nu_{SN} \lesssim 0.2 \text{ yr}^{-1}$, and $L_{SN} \simeq L_K \simeq 10^{43} \text{ erg s}^{-1}$), and smaller than ($\lesssim 20 \%$) the observed bolometric luminosities of the sources ($L_{bol} \simeq L_{FIR} \gtrsim 10^{11} L_\odot$ for typical LIGs and Sey galaxies). In fact, we will see later (§3) that only a fraction of this SN energy rate is actually necessary to power the outflow. The remaining SN energy will be possibly lost in radiation, as indicated by recent numerical studies (Thornton, Gaudletz, Janka, & Steinmetz 1998)

3. Modelling the Outflow

3.1. Disk Parameters

In the previous section, we have assumed that a rotating nuclear disk of gas is produced during the SB evolution and have evaluated the amount of *kinetic* energy that such a system is able to inject into the ambient medium. Now, before deriving the conditions for outflow acceleration and collimation from the surface of the disk, we must explore fiducial physical parameters for the disk. In earlier studies of SB-driven, wide-angled super-winds (e.g., HAM90), global physical conditions were derived for the central regions (radius $\lesssim 1$ kpc) of SB galaxies. Here, since we are focusing on the inner regions (located within few tens of pcs radii from the nucleus) of systems with potential coexistence of a SB-AGN activity, we take the observed data of the nuclear disk of the LIG prototype, Arp 220 (SYB97). High-resolution CO imaging and dust continuum emission of the nuclear region of Arp 220 has revealed the presence of a rotating disk feature of molecular gas with mean density $n_{H_2} \simeq 2 \times 10^4 \text{ cm}^{-3}$, a radius $\simeq 250$ pc, and two stellar nuclei at ~ 235 pc from the center. Most of the disk mass resides in the molecular gas, rather than in the stars, and most of the molecular gas ($M_d \sim 5.4 \times 10^9 M_\odot$) is concentrated in a very thin disk within the double nuclei distance from the center. The fact that the bulk of the molecular gas has relaxed into a disk with large mass of gas concentrated interior to the double stellar nuclei is consistent with a scenario in which the gas in merging systems settles into the center faster than the stellar components of the SB. The mass in the Arp 220 disk is approximately distributed according to the empirical relation (see Fig. 8 of SYB97)

$$M_r \simeq 5 \times 10^9 M_\odot \left(\frac{r}{200 \text{ pc}} \right)^\beta \quad (5)$$

with $\beta \simeq 1.66$ (for $r \lesssim 200$ pc).

The rotation velocity is related to the mass distribution through the simple spherical

approximation $v_{rot} = (GM_r/r)^{1/2}$.

For a self-gravitating disk, the effective thickness (H) of the gas can be obtained from the standard condition of hydrostatic equilibrium (Spitzer 1942), $H = \left(\frac{0.233\sigma^2}{G\Sigma}\right)$, where $\sigma \simeq 90 \text{ km s}^{-1}$ is the vertical dispersion velocity (SYB97), and $\Sigma = M_d/A_d = 2H\rho_d$ is the mass surface density of the disk. Within a radius $r \simeq 200 \text{ pc}$ and $M_d \simeq 5 \times 10^9 M_\odot$, $\Sigma \simeq 3 \times 10^4 M_\odot \text{ pc}^{-2}$ for Arp 220, so that

$$H \simeq 15 \text{ pc} \left(\frac{\sigma}{90 \text{ km s}^{-1}}\right)^2 \left(\frac{\Sigma}{3 \times 10^4 M_\odot \text{ pc}^{-2}}\right)^{-1}. \quad (6)$$

This implies a thin disk at $r \simeq 200 \text{ pc}$ with $H/r \simeq 0.075$. In the inner region it can be thicker however.

The stability of the gas disk to axi-symmetric perturbations has been examined by SYB97 who have used the Toomre (1964) criterion $Q \simeq \sigma k/(\pi G \Sigma)$ (where $k \simeq 2\Omega$ is the epicyclic frequency). The fact that $Q \lesssim 1$ within $r \simeq 400 \text{ pc}$ implies that the gas disk is unstable and will develop large density enhancements, ultimately leading to the formation of massive star clusters in the disk. This result is consistent with our previous assumption of a disk originated in a SB environment also processing star formation.

The extreme gas densities in the central disk of Arp 220 ($n_d = n_{H_2} \simeq 2 \times 10^4 \text{ cm}^{-3}$) will result in high accretion rates. Assuming a viscous transport of angular momentum, SYB97 have used the Pringle (1981) treatment to estimate the rate of gas accretion as a function of the disk radius. For the typical Arp 220 parameters, they found accretion rates $\dot{M}_d \simeq 100 M_\odot \text{ yr}^{-1}$ peaking at $\sim 250 \text{ pc}$ (which corresponds to a kinematic viscosity $\nu \simeq 3 \times 10^{27} \text{ cm}^2 \text{ s}^{-1}$).

We now must check whether the above accretion rate $\dot{M}_d$ is consistent with the range of temperatures and pressures estimated for the disk. Accretion through the disk implies that there is dissipation of energy into the disk. For a self-luminous disk the temperature

as a function of the radius is $T_r^4 = \frac{3GM_d\dot{M}_d}{8\pi\sigma_{SB}r^3}$, where σ_{SB} is the Stefan-Boltzmann constant.

Using the accretion rate mentioned above we estimate

$$T_d \simeq 14 \text{ K} \left(\frac{M_d}{5 \times 10^9 M_\odot} \right)^{1/4} \left(\frac{\dot{M}_d}{100 M_\odot \text{ yr}^{-1}} \right)^{1/4} \left(\frac{r}{200 \text{ pc}} \right)^{-3/4} \quad (7)$$

which is consistent with the temperatures estimated from the OH observations in Arp 220 ($\simeq 17\text{--}21$ K; SYB97).

The sound speed at this radius would be $v_s \sim 0.4 (T/14 \text{ K})^{1/2} \text{ km s}^{-1}$, and the disk pressure

$$P_d \simeq 4 \times 10^{-11} \text{ dy cm}^{-2} \left(\frac{n_d}{2 \times 10^4 \text{ cm}^{-3}} \right) \left(\frac{T}{14 \text{ K}} \right). \quad (8)$$

We note, however, that in our SB/disk model, massive star formation may still be occurring after the formation of the disk and even in the disk itself. In such a case, one would expect the presence of an ionized, warmer gas component produced in interactions between the ejected matter in the SB (e.g., HAM90). A disk with a dominant warm gas component (with $T_d \simeq 10^4$ K) would have a higher pressure $P_d \simeq 10^{-8} \text{ dy cm}^{-2}$. In any case, no matter whether the inner disk is cold (i.e., presently not undergoing massive star formation) or warm, the magneto-centrifugal mechanism for outflow production that will be discussed below is applicable to both.

3.2. Outflow Conditions

Previous SB-driven modeling of wide-angled super-winds (e.g., Tomisaka & Ikeuchi 1988, De Young & Gallagher 1990, Suchkov et al. 1994) have shown that the multiple action of stellar winds and SN explosions may lead to the formation of a shell of shocked disk material which, after blowout, is dragged to the halo, where it forms the walls of a bi-conical cavity. This would, ultimately, create conditions for a free directed escape of hot gas at the later, more vigorous SN dominated phase of the SB. Formulation derived by Mac

Low & McCray (1988) and Mac Low, McCray, & Norman (1989) gives a simple condition for a SB-driven outflow to blow out of the atmosphere above the disk. The rate of energy released by the SN, L_K (eq. 4), must satisfy

$$L_K > 3 \times 10^5 L_\odot \left(\frac{n_d}{2 \times 10^4 \text{ cm}^{-3}} \right)^{-1/2} \left(\frac{H}{15 \text{ pc}} \right)^2 \left(\frac{P_d}{10^{-8} \text{ dy cm}^{-2}} \right)^{3/2} \quad (9)$$

which is a condition more than satisfied by the candidate systems, which have $L_K \simeq (5.3 \times 10^7 - 2 \times 10^{10}) L_\odot$ (eq. 4). We note that this condition still holds for more compact, denser disks with, for example, $H \sim 1 \text{ pc}$, $n_d \lesssim 10^8 \text{ cm}^{-3}$, and $P_d < 10^{-3} \text{ dy cm}^{-2}$.

The results above suggest that nuclear SBs may satisfy the conditions for driving winds out of the galactic core of the Seyferts and LIGs, but, how to explain the collimated morphology of the observed outflows that we are investigating and their acceleration? One could invoke a thick gas torus surrounding the core to collimate the "wind" in a similar way to that proposed in standard models for AGN (see, e.g., Antonucci 1993). As mentioned earlier, hydrodynamic models for SB super-winds predict the formation of a cavity that provides a natural channel for vertical escape of the gas ejected by the SN. Recently, Tenorio-Tagle & Muñoz-Tuñón (1997), have explored hydrodynamic solutions in which the rapid fall of disk matter towards the center develops a shock front in the inner region of the disk, and the material crossing it experiences a splash and is ejected vertically to the lower pressure zones above the disk, thus forming a (poorly collimated) kpc-long cone. All these models seem to be able to provide appropriate conditions for the creation of the observed, poor-collimated super-winds emerging from the kpc-scale galactic disks of SB galaxies. Here, in our investigation of the dynamics of nuclear SB/disks, we examine another potential mechanism in which efficient collimation and outflow acceleration is provided by magnetic fields anchored into the core disk of the SB – an assumption that is supported by the observation of magnetic fields in the outflows of the Seyferts ($B \simeq 3.5 \times 10^{-7} - 1.6 \times 10^{-6}$

G at few kpc above the disk; Colbert et al. 1996). Such mechanism could also, in principle, be operating in the inner (possibly unobservable) regions of wide-angled super-winds of normal SB galaxies.

In a magnetized gas under low ionization fraction condition, for an outflow to be launched the neutral component must be well coupled with the ions (e.g. Wardle & Königl 1993). This coupling is described by η , the ratio of the dynamical time to the mean collision time of a neutral atom in a diluted sea of ions (e.g. Wardle & Königl, Shu et al. 1994, 1995), $\eta = \frac{\gamma_i \rho_i r}{v_{rot}}$, where ρ_i is the density of ions and v_{rot} is the rotational velocity of a particle in a circular orbit. When $\eta \gg 1$ the neutrals move with the ions and so can be described as well coupled to the magnetic field. Here $\gamma_i \sim 3 \times 10^{13} \text{ cm}^3 \text{ g}^{-1} \text{ s}^{-1}$ is the drag coefficient between ions and neutrals and ρ_i , the density of ions, is the product of the atomic mass of the ions, the ionization fraction and the total density. We can then estimate

$$\eta > 5.6 \times 10^2 \left(\frac{n_e/n_H}{10^{-5}} \right) \left(\frac{\rho_d}{10^{-19} \text{ g cm}^{-3}} \right) \left(\frac{M_d}{5 \times 10^9 M_\odot} \right)^{-1/2} \left(\frac{r}{200 \text{ pc}} \right)^{3/2} \quad (10)$$

For clouds with temperatures, abundances and column densities not much different from those in the disk examined here, Friaça & Jafelice (1998) have found that the ionization state is roughly inversely correlated with the density. From their curves we have estimated the corresponding value of $n_e/n_H \simeq 10^{-5} - 10^{-2}$ for our density and temperature ranges. We conclude that over the whole disk $\eta \gg 1$ and the neutrals can be said to be well coupled to the ions. However this does not mean that the particles are necessarily well coupled to the magnetic field.

The magneto-centrifugal wind models can operate where the charged particles are sufficiently coupled to the field lines that they are flung out centrifugally from the outer layers of the disk (for a favorable field geometry; see below). This occurs when the magnetic Reynold's number is $Re_M \gg 1$. We can estimate the magnetic Reynolds number as $Re_M = \frac{v_{rot} H}{\nu_M}$, where ν_M is the magnetic resistivity, $\nu_M = c^2/(4\pi\sigma_e)$, and σ_e is

the electric conductivity determined from collisions between electrons and H_2 molecules, $\sigma_e \simeq \frac{n_e e^2}{n_{H_2} m_e \langle \sigma_c v \rangle} \simeq 1.1 \times 10^{16} (n_e/n_H) \text{ s}^{-1}$ (where σ_c is the cross section for $e - H_2$ collisions, v is the relative velocity between them, and $\langle \sigma_c v \rangle$ is the Maxwellian mean of the product; Hayashi 1981, Takata & Stevenson 1996). Using the Arp disk parameters we estimate

$$Re_M > 2.3 \times 10^{18} \left(\frac{M_d}{5 \times 10^9 M_\odot} \right)^{1/2} \left(\frac{H}{15 \text{ pc}} \right) \left(\frac{n_e/n_H}{10^{-5}} \right) \left(\frac{r}{200 \text{ pc}} \right)^{-1/2} \quad (11)$$

so that we can assume that the magnetic field is well coupled to the gas.

3.3. Magneto-centrifugally Driven Outflows

Collimated winds may be produced by a rotating disk with a magnetic field ($\mathbf{B}$) anchored into it (see, e.g., Spruit 1996 and references therein). The magnetic field and disk structure will determine how much mass is launched out of the disk.

As remarked before, we here assume that the field is produced due to magnetic flux capture from the environment during the formation of the disk in the SB, followed by advection and compression by the accretion process. Poloidal flux capture in this way cannot be destroyed by local processes in the disk, it can only escape by diffusing radially outward (e.g., Spruit 1996). It is out of the scope of this work to formulate a theory for (turbulent) diffusion in the accretion disk, so that we cannot reliably predict the distribution of the magnetic field. However, since all the energy densities are expected to increase towards the center, it is reasonable to assume that the field will increase inward as well and form an opening field geometry. As argued by Spruit, Stehle, & Papaloizou (1995), fields with such a configuration can be quite strong, with magnetic energy densities exceeding the gas pressure, which must make them ideal for magnetic outflow production.

Just above the disk, the gas density is typically low enough, so that the magnetic

energy density is large compared with the thermal and rotational energies. The field in this region may be thus assumed to be force-free ($\mathbf{J} \times \mathbf{B} = \mathbf{0}$). The poloidal magnetic field lines that raise from the disk are somewhat like rigid wires that control the flow. The gas of the disk surface will be forced to corotate with the field lines raising out of the disk. For a sufficiently cold disk, the solely forces acting on a parcel of gas will be the gravitational force and the centrifugal force due to rotation. Along the field line above the disk, the centrifugal force will increase with the distance from the axis and the effective gravity decrease. When the centrifugal force component along the line exceeds that of gravity, the gas tied to the field line is accelerated outward. This outward centrifugal acceleration continues up to the Alfvén point [the location where the outflow poloidal speed reaches the Alfvén speed, $v = v_A = B^2/(4\pi\rho)^{1/2}$, and stops when the flow attains the fast magnetosonic point. At that point, the field is no longer strong enough to enforce corotation. Beyond the Alfvén surface ($\rho v^2 > B^2/8\pi$), the inertia of the gas causes it to lag behind the rotation of the field line, and the field winds up thus developing a strong azimuthal component (B_ϕ). The magnetic stress force associated to the azimuthal field [$-B_\phi^2/(4\pi r)$, where r is the radial distance from the disk axis] will provide the flow collimation (see e.g., PP92, Spruit 1996).

A self-gravitating accretion disk with an accretion rate $\dot{M}_d$ releases a known amount of gravitational binding energy of gas as it spirals inward to the center. The presence of viscous stress causes the gravitational binding energy to transform into heating and the angular momentum is radially transported outward by turbulent stress. In the presence of magnetic fields, however, part of the gravitational energy released in the accretion is converted into mechanical energy of a wind. In addition, the angular momentum is efficiently transported by the wind through the disk surface, and the wind mass-loss rate $\dot{M}_w$ has only to be a tiny fraction of $\dot{M}_d$ (see, e.g., PP92 and references therein). Using the disk angular momentum equation one can show that as long as the magnetic pressure is comparable or larger than the gas thermal pressure, the magnetic torque will completely dominate the viscous torque

(e.g. PP92). In fact, the ratio of the magnetic to the viscous torque is

$$\left(\frac{\tau_w}{\tau_{vis}}\right) = \left(\frac{B_z^2}{4\pi P_d}\right) \left(\frac{r_A}{\alpha_{SS}H}\right) \quad (12)$$

where $\alpha_{SS} < 1$ is the Shakura-Sunyaev viscous prescription (e.g., Frank, King, & Raine 1992), and r_A is the Alfvén radius. Thus, since $r_A/\alpha_{SS}H > 1$, and the magnetic pressure in the disk ($B^2/8\pi \gtrsim 2.5 \times 10^{-8} \text{ dy cm}^{-2}$; see below) is effectively of the order or larger than the thermal pressure (Eq. 8), we expect that the angular momentum loss through the wind will dominate.

In magneto-centrifugal disk outflows, the angular momentum of the wind is tied to that of the disk, and so the wind flux is directly related to the accretion rate, $\dot{M}_d$, through the disk. To investigate specific conditions under which collimated winds could be generated in the SB/disk scenario drawn in the previous paragraphs, we here follow the non-self-similar, steady-state, magneto-centrifugally driven wind models developed by PP92. In their solutions, the angular velocity of the disk is assumed to be a power law of the magnetic flux, $\Omega \propto \psi^{-\alpha}$. They have found that for a power law index $\alpha = 3$, the outflow carries a constant current intensity everywhere and is free of the divergence found in self-similar models (e.g., Blandford & Payne 1982). We here adopt this relation. Now, according to Eq. 5, the angular velocity scales with the disk radius as

$$\Omega = \frac{(GM_r)^{1/2}}{r^{3/2}} \propto r^{(\beta-3)/2} \propto r^{-0.67}. \quad (13)$$

Therefore, the magnetic flux will also be a power law of the radius,

$$\psi \propto r^{\frac{(3-\beta)}{2\alpha}} \propto r^{0.223}. \quad (14)$$

This, in turn, implies the following relation for the poloidal magnetic field in the disk as a function of the radius (PP92), $B_r \propto r^{-\delta}$, with $\delta = 2$, or

$$B_r = B_\infty \left(\frac{r_\infty}{r}\right)^\delta \quad (15)$$

where $B_\infty \simeq 10^{-5}$ G is the average value inferred from the observations of the Sey outflows at the kpc-scale, $r_\infty = 1000$ pc.

In these models, the ratio of the radius r to the Alfvén radius, r_A , $\frac{1}{3}(r/r_A)^3$, can be determined by the parameter (PP92)

$$\epsilon \equiv \frac{\dot{M}_d \sqrt{G M_r}}{3 B_r^2 r^{5/2}} \quad (16)$$

For a wind to be produced the field lines must be more than 20° from normal to the disk and $\epsilon \leq 1$. Assuming $\dot{M}_d$ constant, the relation above for B_r (Eq. 15) and Eq. 5 for the disk mass result

$$\epsilon \propto r^{\frac{(\beta-5)}{2}+2\delta} \propto r^{2.33}. \quad (17)$$

Using these relations and the physical conditions in the Arp 220 disk (with constant $\dot{M}_d = 100 M_\odot \text{ yr}^{-1}$), we can determine a maximum radius r_e for which $\epsilon(r_e) = 1$,

$$r_e \simeq 127 \text{ pc} \left(\frac{\dot{M}_d}{100 M_\odot \text{ yr}^{-1}} \right)^{-0.43} \left(\frac{B_\infty}{10^{-5} \text{ G}} \right)^{0.86} \left(\frac{r_\infty}{1 \text{ kpc}} \right)^{1.72} \quad (18)$$

This value determines a characteristic maximum radius for the outflow zone in the disk.

We note, however, that it is sensitive to the adopted value of the magnetic field on the large scales B_∞ which is only barely known. (Notice that Eq. 15 implies $B(r_e) \simeq 6 \times 10^{-4}$ G on the disk surface at r_e .)

For these models the mass flux in the wind, $\dot{M}_w$, is related to the accretion rate through the disk by (PP92)

$$\dot{M}_w = f \dot{M}_d \quad (19)$$

where

$$f \approx \left(\frac{r}{r_A} \right)^2. \quad (20)$$

HAM90 have estimated values for $\dot{M}_w$ for wide-angled super-winds observed in far-infrared galaxies ranging from a few $M_\odot \text{ yr}^{-1}$ in the low-power sources (e.g., M82), up

to $\simeq 50 M_\odot \text{ yr}^{-1}$ in powerful sources (like Arp 220 and Mrk 273). These values correspond to the total amount of mass loss rate integrated over the whole radius of those winds. In the inner, few tens of pc scales, $\dot{M}_w$ is expected to be smaller. In fact, in our model, $\dot{M}_w$ approximately scales as $\dot{M}_w \propto \psi^2$ (eq. 6.7 of PP92), or according to Eq. 14 above, $\dot{M}_w \propto r^{0.45}$. Therefore, the value $\dot{M}_w \simeq 50 M_\odot \text{ yr}^{-1}$ for the Arp 220 outflow will correspond to disk radius $\gg 100 \text{ pc}$. We can then assume that in the inner few pcs $\dot{M}_w \simeq 10 M_\odot \text{ yr}^{-1}$ and take $\dot{M}_w/\dot{M}_d \simeq 0.1$. This value is compatible with estimates for YSOs and AGN outflow models (see, e.g. PP92 and references therein). Substituting this value into Eqs. 19, and 20 above, we can then estimate a fiducial value for the ratio $r_A/r_i \simeq (\dot{M}_w/\dot{M}_d)^{-1/2} \simeq 3.2$ at a fiducial radius r_i , and according to Eq. 16,

$$\epsilon_i \simeq \frac{1}{3} \left(\frac{r_i}{r_A} \right)^3 \simeq 1.1 \times 10^{-2} \quad (21)$$

Thus substituting this value into Eq. 17,

$$r_i = r_e \left(\frac{\epsilon_i}{\epsilon_e} \right)^{0.43} \simeq 18 \text{ pc}, \quad (22)$$

or (according to Eq. 16)

$$\epsilon_i \sim 1.1 \times 10^{-2} \left(\frac{\dot{M}_d}{100 M_\odot \text{ yr}^{-1}} \right) \left(\frac{M_{r_i}}{9.2 \times 10^7 M_\odot} \right)^{1/2} \left(\frac{B_{r_i}}{3.1 \times 10^{-2} \text{ G}} \right)^{-2} \left(\frac{r_i}{18 \text{ pc}} \right)^{-5/2}. \quad (23)$$

The terminal speed of the outflow can be evaluated from the solution of the Bernoulli equation for the flow in the asymptotic domain (e.g., PP92, eq. 6.6)

$$v_\infty(r) \simeq \frac{(2\lambda)^{1/2}}{3 \epsilon(r_e)} (\Omega_e r_e) \psi^{-2} \propto r^{-0.45} \quad (24)$$

where $(\Omega_e r_e)$ is the rotation velocity in the disk at the outer outflow radius r_e , and λ is a constant everywhere in the wind that can be estimated from Eq. 6.7 of PP92, and Eq. 14 above

$$\lambda \sim \frac{4.5}{2} \epsilon(r_e)^2 \frac{\dot{M}_d}{\dot{M}_w(r_e) - \dot{M}_w(r_i)} \left[1 - \left(\frac{r_i}{r_e} \right)^{0.45} \right] \simeq 9.35 \quad (25)$$

where we have used the approximate scaling law for $\dot{M}_w(r) \propto \psi^2 \propto r^{0.45}$. Substituting λ into Eq. 24, we find the expected terminal velocity at the radius r_i at which most of the wind originates

$$v_\infty(r_i) \simeq 973 \text{ km s}^{-1} \left(\frac{\Omega_e r_e}{282 \text{ km s}^{-1}} \right) \left(\frac{r_i/r_e}{0.14} \right)^{-0.45} \quad (26)$$

and $v_\infty(r_e) \simeq 0.42 v_\infty(r_i) \simeq 408 \text{ km s}^{-1}$. These estimates are consistent with the detection of emission lines typically with widths $\sim 200 - 600 \text{ km s}^{-1}$ in LIGs. They indicate the presence of a high speed pre-shocked wind gas that was possibly decelerated to these velocities. In the case of Arp 220, the observation of line widths declining from $\sim 700 \text{ km s}^{-1}$ near the axis of the outflow to $\sim 300 \text{ km s}^{-1}$ at the outer radius are also consistent with the prediction of our model of a decreasing outflow velocity with radius. On the other hand, the estimates above are higher than the observed outflow speeds in the Seyfert outflows ($v_{\infty,obs} \simeq 25 - 150 \text{ km s}^{-1}$; Condon et al. 1996). Nonetheless, according to Eq. 24, these observations can be fitted by a more compact disk (like that of Fig. 1, for instance) with a somewhat larger ratio $\dot{M}_w/\dot{M}_d$ (Eq. 21). Eqs. 21, 22, 24 and 25 indicate that disks with $r_e \lesssim 10 \text{ pc}$ and $\dot{M}_w(r_i)/\dot{M}_d \gtrsim 0.1$ will provide v_∞ in better agreement with the observed Sey outflows.

In summary, the results above indicate that the observed terminal outflow velocities are consistent with an outflow generation in the described SB-disk scenario with wind mass loss to disk accretion rate ratios $\gtrsim 0.1$.

One of the attractions of a magnetically driven outflow model is that it is able not only to produce and collimate the outflow, but also to efficiently extract angular momentum from the disk, allowing it to collapse (e.g. Blandford 1976, PP92). Just a small fraction of the infalling mass, like that estimated above, is required to be carried out by the outflow for it to take out most of the angular momentum of the system. Further, in the outflow model investigated here, it is easy to show that the total kinetic power carried by the

wind is exactly one-half of the gravitational energy released in the accretion disk, i.e., $L_w = \frac{1}{2} L_{acc} = \frac{1}{2} \frac{GM_{re}\dot{M}_d}{r_e} \left[1 - \left(\frac{r_i}{r_e} \right)^{0.45} \right]$, where M_{re} is the disk mass within r_e . Again, using the Arp 220 parametrization, we find that

$$L_w(r_i) \simeq 1.5 \times 10^{42} \text{ergs}^{-1} \left(\frac{\dot{M}_d}{100 M_\odot \text{yr}^{-1}} \right) \left(\frac{M_{re}}{2.3 \times 10^9 M_\odot} \right) \left(\frac{r_e}{127 \text{pc}} \right)^{-1} \left[1 - \left(\frac{r_i/r_e}{0.14} \right)^{0.45} \right]. \quad (27)$$

This can be compared with the SN energy rate estimated for the Arp 220 disk, $L_K \simeq 7 \times 10^{43} \text{erg s}^{-1}$ (Eq. 4), which results $L_w/L_K \simeq 2 \times 10^{-2}$, so that the total kinetic power carried by the magneto-centrifugally driven outflow is much smaller than the available SN energy rate predicted by the SB models.

The fact that the open field lines must have an inclination angle with respect to the disk not larger than $\theta = 70^\circ$ (PP92) results that the height z_A of the Alfvén surface is $z_A < r_A \tan \theta$, or

$$z_A < 150 \text{pc} \left(\frac{r_A/r_i}{3} \right) \left(\frac{r_i}{18 \text{pc}} \right). \quad (28)$$

We can also estimate the outflow number density at the Alfvén surface (and beyond) from the condition that $n_A \propto \psi^2$ (PP92), and according to Eq. 14, $n_A \propto r^{\frac{3-\beta}{\alpha}} \propto r^{0.45}$. Using the number density at the base of the disk, which was estimated from the Arp 220 disk, $n(r_i) \simeq 3 \times 10^4 \text{cm}^{-3}$, the relation above implies that

$$n_A \simeq 5 \times 10^4 \text{cm}^{-3} \left(\frac{r_A/r_i}{3} \right)^{0.45} \left(\frac{r_i}{18 \text{pc}} \right)^{0.45}. \quad (29)$$

According to PP92 models, the flow will recollimate beyond the fast-magnetosonic (fm) point. The Mach number at this point is

$$M_{fm} = (2^{3/2} \lambda^{1/2} - 3)^{1/2} \quad (30)$$

Substituting the value of λ given by Eq. (25) we find $M_{fm} \simeq 2.4$. Thus flows achieving

modest fm-Mach number ~ 2 must recollimate and form magnetic focal points beyond about a height $z_A \sim 150$ pc.

4. Conclusions and Discussion

SBs with a disk core formed by infalling gas produced during star formation in a rotating stellar cluster, seem to be able to provide the amount of energy required to drive the weak, directed outflows of Seyferts galaxies and LIGs. In the latter source class, for example, there seems to be strong observational support for the presence of nuclear SBs (e.g., SLL98). We have found that massive nuclear SBs with cluster masses $M_c \sim 10^9 - 10^{10} M_\odot$, disk masses $M_d \sim 10^8 - 10^9 M_\odot$, and supernova rates $\nu_{SN} \simeq 5 \times 10^{-3} - 2 \text{ yr}^{-1}$ (which are consistent with the inferred ν_{SN} from the observed non-thermal radio power) may inject kinetic energies $\sim 10^{41} - 10^{43} \text{ erg s}^{-1}$ which are high enough to blow out directed winds from the accreting disk surface (§2), within the SB lifetimes.

In our models, the acceleration and collimation of the outflow is provided by moderate magnetic fields anchored into the rotating SB-disk. Based on empirical conditions derived from the disk of the prototype LIG, Arp 220, and the magnetized outflows of Sey galaxies and LIGs, we have found that magnetic fields with intensities $\gtrsim 8 \times 10^{-4} - 5 \times 10^{-2} \text{ G}$ are able to accelerate the flow up to the observed terminal velocities ($\lesssim \text{few } 100 \text{ km s}^{-1}$ in the case of the Seyferts, and $\sim 400 - 950 \text{ km s}^{-1}$ in the case of the LIGs), with wind mass loss to disk accretion rate ratios $\dot{M}_w/\dot{M}_d \gtrsim 0.1$ (where $\dot{M}_d \sim 100 M_\odot \text{ yr}^{-1}$). The outflow carries only a small fraction (few percent) of the available SN kinetic energy rate, and is produced within a wind zone in the disk of radius $\lesssim 100 \text{ pc}$ in the case of the LIGs, and $\lesssim 10 \text{ pc}$ in the case of the Seyfert outflows (§3). The model also predicts a decay of the terminal velocity with radius which is compatible with observations.

The formation of an outflow in the proposed SB-magnetized disk scenario depends on a series of conditions which are evolutionary linked, such as, the initial amount of angular momentum and ellipticity in the central stellar cluster that will provide the formation of the rotating disk of gas from the SB; the development of an appropriate magnetic field geometry with open lines during disk collapse; and the ionization state of the gas which must be large enough to couple the magnetic field to the gas. SB-driven outflows are observed not only in IR luminous nuclear SB systems, but also in other galaxy types, as for example, blue compacts and irregular dwarf galaxies like NGC 1569 (e.g., Greggio et al. 1998). The collimation-acceleration mechanism here described could, in principle, also apply to the inner regions of the wide-angled super-winds observed in these systems. They could also anchor a nuclear magnetized disk generated during the SB if the initial conditions for the SB were not very much different from those described here. However, in the case of the amorphous dwarf systems like NGC 1569, which is actually a post-SB system, it is not unlikely that the SB-driven large-scale wind had stripped the dwarf of most of its gas before the gas had time to partially collect into the center and develop a nuclear rotating disk and a more collimated, coherent outflow portion. In forthcoming work, we aim to explore self-consistent numerical models to examine the dynamical evolution of outflows produced by a rotating magnetized disk originated in different SB environments and determine their range of applicability. For that purpose, we will employ a modified version of a 3-D numerical SPH code originally designed for supersonic jet studies (e.g., de Gouveia Dal Pino & Benz 1993, de Gouveia Dal Pino & Birkinshaw 1996, Cerqueira & de Gouveia Dal Pino 1998).

Pure hydrodynamic models (e.g., Suchkov et al. 1994) make specific predictions for the observed abundance pattern in the SB-driven super-winds. In those models, the material at the wall of the vertical cavity, for example, is mostly composed of unprocessed ISM gas that has been shock-heated by the emerging outflow, while the enriched material that is

ejected by the SN should fill the interior of the cavity. Similarly, as a collimated magnetized outflow impinges supersonically the ambient medium, it must create a bow shock pattern at the outer edge (see, e.g., Cerqueira, de Gouveia Dal Pino, & Herant 1997) where most of the shocked material that produces the emission lines is made of ISM gas with no significant abundance enrichment. The observed optical emission lines should then trace normal abundances just like predicted in hydrodynamical models. The hotter, enriched material that is ejected by the SN should emerge mainly along with the collimated outflow.

We note that in our disk model the magnetic pressure in the disk may exceed the gas pressure. This may cause the disk to become unstable to magnetic buoyancy. In such a case, the magnetic field lines may arise in loops from the disk surface and undergo reconnection. This could in turn provide the formation of hot filamentary and clumpy structure on the disk corona. It is interesting to note that a *patchy* disk corona is required by the X-ray and γ -ray observations of the nuclear region of Sey galaxies in order to explain their complex high-energy spectra (e.g., Johnson et al. 1997).

Finally, we should inquire about the fate of the SB disk (see also GT97). We have seen that it may rapidly become self-gravitating (§2.2). As a consequence, it is expected to suffer fragmentation and star formation. Either, the disk may become bar unstable, a condition which is met if $\delta_b = T_{rot}/W > 0.14$, where T_{rot} is the rotational energy and W is the gravitational potential energy of the disk (e.g. Shlosman et al. 1989). Our SB disks result $\delta_b = 0.5$, so that they can, in principle, become unstable to bar formation and undergo fission into two components which eventually may collapse to a supermassive BH (Begelman, Blandford, & Rees 1980). Based on the fact that the formation of BHs seems to be plausible in a SB, we can speculate that both, the standard BH and the SB mechanisms may be occurring in different stages of the AGN evolution. In this evolutionary scenario, the stronger more collimated jets of radio loud QSOs, for example, could be produced in

the higher-level activity phase of the BH accreting the remains of the disk material. This hypothesis is examined elsewhere.

This work has been partially supported by grants of the Brazilian agencies FAPESP and CNPq. E.M.G.D.P. acknowledges the kind hospitality of the staff of the Abdus Salam International Centre for Theoretical Physics, in Trieste, where this work was accomplished. We are also indebted to the referee for his comments and suggestions that we believe have helped to improve this work.

REFERENCES

- Antonucci, R. R. J. 1993, ARA&A, 31, 473
- radio jets
- Blandford, R. D., & Payne, D. G. 1982, MNRAS, 199, 883
- Begelman, M., Blandford, M. C. & Rees, M.J. 1980, Nature, 287, 307
- Begelman, M., & Rees, M.J. 1978, ApJ
- Bryant, P.M., & Scoville, N.Z. 1996, ApJ, 457, 678
- Cerqueira, A.H.& de Gouveia Dal Pino, E.M., 1998, ApJ (in press)
- Cerqueira, A.H., de Gouveia Dal Pino, E.M. & Herant, M. 1997, ApJ, 489, L185
- Colbert, E. J. M. et al. 1996, ApJ, 467, 551
- Condon, J. J. & Yin, Q. F. 1990, ApJ, 357, 97
- de Gouveia Dal Pino, E.M. & W. Benz 1993, ApJ, 410, 686
- de Gouveia Dal Pino, E.M. & Birkinshaw, M. 1996 ApJ, 471, 832
- de Gouveia Dal Pino, E. M. & Medina Tanco, G. A. 1997, in Starburst Activity in Galaxies,
eds. P. Franco, G. Tenorio-Tagle & R. Terlevich, RMAA, series confs., 6, 235 (GT97)
- De Young, , D. & Gallagher, J. 1990, ApJ, 356, L15
- Frank, J., King, A., & Raine, D. 1992, in Accretion Power in Astrophysics, Cambridge
Univ. Press (2nd ed.)
- Field, J.B. 1964, ApJ, 140, 1434

- Friaca, A.C.S., & Jafelice, L.C. 1998, MNRAS (in press)
- Gregio, L. et al. 1998, astro-ph/9803326
- Hartmann, L., & MacGregor, K. B. 1982, ApJ, 259, 180
- Hayashi, C. 1981, Prog. Theor. Phys. Suppl., 70, 35
- Heckman, T. M., Armus, L. & Miley, G. K. 1990, ApJ (Suppl.), 74, 833 (HAM90)
- Heckman, T.M., Lehrnet, M.D. & Armus, L. 1993, in The Environment and Evolution of Galaxies, ed. J.M. Shull & H.A. Thronson, Jr. (Dordrecht: Kluwer), 455
- Heyvaerts, J., & Norman, C. 1989, ApJ, 347, 1055
- Jogee, S., Kenney, J.D., & Smith, B.J. 1998, ApJ, 494, L
- Johnson, W.N. et al. 1997, in Proc. of the IV Compton Symp., Eds. Dermer, C.D., Strickman, M.S., & Kurfess, J.D., AIP Conf. ser., part 1, 410, 283
- Kennicutt, R.C. et al. 1987, AJ, 93, 1011
- Lo, K.Y. et al., 1987, ApJ., 312, 574
- Mac Low, M., & McCray, R. 1988, ApJ, 324, 776
- Mac Low, M., & McCray, R., & Norman, M. 1989, 337, 141
- Mestel, L. 1968, MNRAS, 138, 359
- Okamoto, I. 1998, MNRAS (in press)
- Pelletier, G. & Pudritz, R. E. 1992, ApJ, 394, 117
- Perry, J. J. & Dyson, J. E. 1985, MNRAS, 235, 665

- Pringle, J.E. 1981, ARA&A, 19,137
- Pudritz, R. E., & Norman, C. A. 1983, ApJ, 274, 677
- Quillen, A.C. & Trilling, E. 1998, ApJ (in press)
- Rees, M. J. 1984, ARA&A, 22, 471
- Schaller, et al. 1992, AA (*Suppl.*), 96, 26 Press), 233
- Scoville, N.Z. & Norman, C.A. 1988, ApJ, 332, 163
- Scoville, N.Z. & Soifer, B.T. 1991, in Massive Stars in Starbursts, Ed. C. Leitherer et al. (New York: Cambridge Univ.
- Scoville, N.Z., Yun, M.S., & Bryant, P.M. 1997, ApJ, 484, 702
- Shklovskii, I.S. 1960, Sov. Astron. A.J., 4, 885
- Shlosman, I., Frank, J. & Begelman, M. C. 1989, Nature, 338, 45
- Shu, F., Najita, J., Ostriker, E., Wilkin, F., Ruden, S. & Lizano, S. 1994, ApJ, 429, 781
- Shu, F. H., Najita, J., Ostriker, E., & Shang, H. 1995, ApJ, 455, L155
- Spitzer, L. 1942, ApJ, 95, 329
- Spitzer, L. & Saslaw, W. C. 1966, ApJ, 143, 400
- Spruit, H.C. 1996, in *Physical Process in Binary Stars*, Kluwer Dordrecht (NATO/ASI ser.)
- Spruit, H.C., Stehle, H., & Papaloizou, J.C.B. 1995, MNRAS, 275, 1223
- Suchkov, A., Balsara, d., Heckman, T., & Leitherer, C. 1994, ApJ, 430, 511

Takata, T., & Stevenson, D. J. 1996, *Icarus*, 123, 404

Tenorio-Tagle, G. & Muñoz-Tuñón, C. 1997, *ApJ*

Terlevich, R. J. et al. 1992, *MNRAS*, 255, 713

Thornton, K., Gaudletz, M., Janka, H.TH., & Steinmetz, M. 1998, *ApJ*, 500, 95

Tomisaka, K. & Ikeuchi, S. 1988, *ApJ*, 330, 695

Wardle, M., & Königl, A. 1993, *ApJ*, 410, 218

Weber, E. J. & Davis, L. 1967, *ApJ*, 148, 217

Williams, R.J.R. & Perry, J.J. 1994, *MNRAS*, 269, 538

Figure Caption

Figure 1. a) Evolution of a stellar cluster and its core disk, with a Salpeter IMF ($M_l = 1M_\odot$, $M_u = 120M_\odot$), $N_c = 3 \times 10^8$, $r_c = 53$ pc, total cluster mass $M_c = 9.5 \times 10^8 M_\odot$, $R_d = 2.6$ pc, total number of stars in the disk $N_d = 9 \times 10^7$, and total stellar mass in the disk $M_d = 1.6 \times 10^8 M_\odot$. Luminosity sources shown: dotted - supernovae in the cluster and disk; solid - rate of energy released by the infalling gas in the disk; dashed - starlight in the cluster and disk; and dot-dashed - the Eddington luminosity limit in the disk. b) The corresponding rates of supernovae production ν_{SN} in the cluster and the disk. For a system with masses and radii about ten times larger (smaller) than those above, the luminosity values and supernova rates increase (decrease) by an order of magnitude.

